# **Chapter 10: Genes** Knowledge organiser **B**1

## Variation

- The differences in characteristics of living things is known as **variation**
- There is a large amount of variation between different species, but within species many more characteristics are shared
- Even though two organisms may look the same, they will always have variation between them

Inherited variation	Environmental variation
<ul> <li>Is anything that comes directly from your parents, anything that you inherit</li> </ul>	<ul> <li>Is any type of variation that is caused by your surroundings</li> </ul>
Examples can include lobe less or lobed	Factors that can cause environmental variation
ear lobes and eye colour	include diet, education and lifestyle

- Environmental factors can also impact inherited factors, for example a poor diet can affect height or your exposure to the sun can affect skin tone
- Characteristics which are inherited and not affected by environmental variation include natural eye colour, blood group and genetic diseases



# The menstrual cycle

۲

- The **menstrual cycle** is the process in which an egg is released from an ovary and leaves through the vagina
- Day 1: blood from the uterus lining leaves through the vagina, which is known as a **period**
- Day 5: the bleeding stops and the uterus lining starts to re-grow
- Day 14: an egg is released from one of the ovaries during ovulation
- If the egg is **fertilised** than the menstrualcycle stops until the baby is born Dav 28 Dav



- Adaptations are characteristics which organisms have developed to best survive in their surroundings
- their genes

• Egg cells and sperm cells are also called **gametes**, and each contains half the genetic information needed to form a complete organism.

### Egg cells

- An egg is released by the ovaries every month
- The egg cell is moved along the oviduct towards the uterus by cilia

- During **gestation** the nutrients from the mother, these are passed through the placenta which is connected to the fetus by the umbilical cord
- Nutrients are passed from the mother to the baby and waste products are passed
- sac which acts as a shock absorber

Make sure you can write definitions for these key terms.

adaptation adolescence amniotic sac cervix cilia egg cell embryo environmental variation fertilisation fetus gamete gestation implantation inherited variation menstrual cycle ovary oviduct ovulation penis period placenta puberty reproductive system scrotum semen sex hormones species sperm cell sperm duct testicles umbilical cord urethra uterus vagina variation

۲



# Adaptations

- Organisms with the best suited adaptations can breed and pass these on
- Those who are not best adapted will die out and not be able to pass on

# Fertilisation, implantation and gestation



#### Sperm cells

Sperm cells are produced in the testicles/testes

Sperm are mixed with nutrients and fluid from the glands to form semen

During sexual intercourse a man will release semen into the vagina (ejaculation)

If a sperm meets the egg fertilisation may happen

The fertilised egg may then **implant** in the uterus lining and form an embryo (ball of cells)

Just a dot 1 week – cells beginning developing fetus needs to specialise 3 mm long 4 weeks – spine and brain 5 forming, heart beating 3 cm long 9 weeks - tiny movements, lips and cheeks sense touch, eyes and ears forming 7 cm long 12 weeks – fetus uses its muscles to kick, suck, swallow, and practise breathing back from the baby to the mother • The baby is protected from bumps to the mother by the **amniotic**